Dicotyledons Aconitum napellus *Cardamine pratensis* Centaurea nigra *Cerastium holostoides Cirsium palustre* Crepis capillaris Daucus carota *Galium verum Hypochaeris radicata Leucanthemum vulgare* Lotus corniculatus Lychnis flos-cocull Malva moschata Origanum vulgare? or Origanum majorana?* Plantago lanceolata Primula veris Primula vulgaris Prunella vulgaris Ranunculus acris Rhinanthus minor Rumex acetosa *Rumex obtusifolius* Sanguisorba minor Stellaria graminea Taraxacum officinale Thymus polytrichus or Thymus vulgaris* Trifolium dubium Trifolium pratense Trifolium repens Vicia sativa

Grasses

Agrostis capillaris Agrostis stolonifers Anthoxanthum odoratum Bromus sp. Cynosurus cristatus Dactylus glomerata Festuca rubra Holcus lanatus Lolium perenne Phleum pratense

Monk's-hood* Cuckooflower Common Knapweed Common Mouse-ear Marsh Thistle Smooth Hawk's-beard Wild Carrot Lady's Bedstraw Cat's-ear Oxeve Daisy Common Bird's-foot-trefoil Ragged-Robin Musk-mallow Wild Marjoram Pot Marjoram **Ribwort Plantain** Cowslip Primrose Selfheal Meadow Buttercup Yellow Rattle Common Sorrel **Broad-leaved Dock** Salad Burnet Lesser Stitchwort Dandelion Wild Thyme Garden Thyme Lesser Trefoil Red Clover White Clover Common Vetch

Common Bent Creeping Bent Sweet Vernal Grass

Crested Dog's tail Cock's-foot Red Fescue Yorkshire Fog Perennial Rye Grass Timothy

The total number of species recorded was 39 including those marked * *Garden species have no place in a wild flower meadow.

Continued on page 2

The flora of the meadow has been under-recorded because recording had to be done from the sidelines to avoid trampling.

Recorded Insects

Bumble Bees and Honey Bes were not as abundant as in 2009. Was this the result of the very cold and prolonged winter?

Butterflies recorded were: Meadow Brown, Ringlet, Small Skipper and Large White

<u>References</u> Stace, Clive (1977). New Flora of the British Isles. CUP

Copyright Survey and records compiled by Ursula Bowen 2010

MANAGEMENT OF THE SHAKESPEARE MEADOW FROM AUTUMN 2010 TO SPRING 2011

The management suggested for autumn 2009 to spring 2010 is still relevant but in view of the very cold and prolonged winter did not take place. These management activities should still be considered.and put into practice.

UB November 2010